

Sandwich classification for classical-like groups over commutative rings

Raimund Preusser

University of Brasilia

- 1 Introduction
- 2 The general linear group $GL_n(R)$
- 3 Sandwich classification for $GL_n(R)$
- 4 The even-dimensional orthogonal group $O_{2n}(R)$
- 5 Sandwich classification for $O_{2n}(R)$
- 6 Sandwich classification for $U_{2n}(R, \Lambda)$
- 7 References

Throughout this talk R denotes an associative, commutative ring with $1 \neq 0$ and $n \geq 3$ a natural number. The following theorem is known as Sandwich Classification Theorem.

Throughout this talk R denotes an associative, commutative ring with $1 \neq 0$ and $n \geq 3$ a natural number. The following theorem is known as Sandwich Classification Theorem.

Theorem (Wilson-Golubchik, 1972/73)

Let H be a subgroup of $GL_n(R)$. Then H is normalized by $E_n(R)$ iff

$$E_n(R, I) \subseteq H \subseteq C_n(R, I)$$

for some ideal I of R .

Throughout this talk R denotes an associative, commutative ring with $1 \neq 0$ and $n \geq 3$ a natural number. The following theorem is known as Sandwich Classification Theorem.

Theorem (Wilson-Golubchik, 1972/73)

Let H be a subgroup of $GL_n(R)$. Then H is normalized by $E_n(R)$ iff

$$E_n(R, I) \subseteq H \subseteq C_n(R, I)$$

for some ideal I of R .

The ideal I in the theorem above is uniquely determined, namely

$$I = I(H) := \{x \in R \mid t_{kl}(x) \in H \text{ for some } k \neq l\}.$$

Let $\sigma \in GL_n(R)$ and set $H := E_n(R)\langle\sigma\rangle$, i.e. H is the smallest subgroup of $GL_n(R)$ which contains σ and is normalized by $E_n(R)$. By the Sandwich Classification Theorem one has $H \subseteq C_n(R, I)$ where $I = I(H)$. That amounts to saying that for any $i \neq j$ and $k \neq l$ the matrices $t_{kl}(\sigma_{ij})$ and $t_{kl}(\sigma_{ii} - \sigma_{jj})$ can be expressed as products of matrices of the form ${}^\epsilon\sigma^{\pm 1}$ where $\epsilon \in E_n(R)$.

Let $\sigma \in GL_n(R)$ and set $H := E_n(R)\langle\sigma\rangle$, i.e. H is the smallest subgroup of $GL_n(R)$ which contains σ and is normalized by $E_n(R)$. By the Sandwich Classification Theorem one has $H \subseteq C_n(R, I)$ where $I = I(H)$. That amounts to saying that for any $i \neq j$ and $k \neq l$ the matrices $t_{kl}(\sigma_{ij})$ and $t_{kl}(\sigma_{ii} - \sigma_{jj})$ can be expressed as products of matrices of the form ${}^\epsilon\sigma^{\pm 1}$ where $\epsilon \in E_n(R)$.

Problem: Find such expressions of the matrices $t_{kl}(\sigma_{ij})$ and $t_{kl}(\sigma_{ii} - \sigma_{jj})$ with a small number of factors.

- 1 Introduction
- 2 The general linear group $GL_n(R)$
- 3 Sandwich classification for $GL_n(R)$
- 4 The even-dimensional orthogonal group $O_{2n}(R)$
- 5 Sandwich classification for $O_{2n}(R)$
- 6 Sandwich classification for $U_{2n}(R, \Lambda)$
- 7 References

Definition

$GL_n(R) := M_n(R)^*$ is called *general linear group*.

Definition

$GL_n(R) := M_n(R)^*$ is called *general linear group*.

Definition

Let $i, j \in \{1, \dots, n\}$ such that $i \neq j$ and $x \in R$. Then $t_{ij}(x) := e + xe^{ij}$ is called an *elementary transvection*. The subgroup of $GL_n(R)$ generated by all elementary transvections is called *elementary subgroup* and is denoted by $E_n(R)$.

Definition

$GL_n(R) := M_n(R)^*$ is called *general linear group*.

Definition

Let $i, j \in \{1, \dots, n\}$ such that $i \neq j$ and $x \in R$. Then $t_{ij}(x) := e + xe^{ij}$ is called an *elementary transvection*. The subgroup of $GL_n(R)$ generated by all elementary transvections is called *elementary subgroup* and is denoted by $E_n(R)$. Let I be an ideal of R . An elementary transvection $t_{ij}(x)$ is called *I -elementary* iff $x \in I$. The subgroup of $GL_n(R)$ generated by all I -elementary transvections is called *preelementary subgroup of level I* and is denoted by $E_n(I)$.

Definition

$GL_n(R) := M_n(R)^*$ is called *general linear group*.

Definition

Let $i, j \in \{1, \dots, n\}$ such that $i \neq j$ and $x \in R$. Then $t_{ij}(x) := e + xe^{ij}$ is called an *elementary transvection*. The subgroup of $GL_n(R)$ generated by all elementary transvections is called *elementary subgroup* and is denoted by $E_n(R)$. Let I be an ideal of R . An elementary transvection $t_{ij}(x)$ is called *I -elementary* iff $x \in I$. The subgroup of $GL_n(R)$ generated by all I -elementary transvections is called *preelementary subgroup of level I* and is denoted by $E_n(I)$. The normal closure of $E_n(I)$ in $E_n(R)$ is called *elementary subgroup of level I* and is denoted by $E_n(R, I)$.

Lemma

The relations

$$t_{ij}(x)t_{ij}(y) = t_{ij}(x + y), \quad (\text{R1})$$

$$[t_{ij}(x), t_{hk}(y)] = e \text{ and} \quad (\text{R2})$$

$$[t_{ij}(x), t_{jk}(y)] = t_{ik}(xy) \quad (\text{R3})$$

hold where $i \neq k, j \neq h$ in (R2) and $i \neq k$ in (R3).

Definition

Let I be an ideal of R . Then the canonical map $R \rightarrow R/I$ induces a group homomorphism $\phi : GL_n(R) \rightarrow GL_n(R/I)$. The preimage of $Center(GL_n(R/I))$ under ϕ is called *full congruence subgroup of level I* and is denoted by $C_n(R, I)$.

Definition

Let I be an ideal of R . Then the canonical map $R \rightarrow R/I$ induces a group homomorphism $\phi : GL_n(R) \rightarrow GL_n(R/I)$. The preimage of $Center(GL_n(R/I))$ under ϕ is called *full congruence subgroup of level I* and is denoted by $C_n(R, I)$.

Remark

Let $\sigma \in GL_n(R)$ and I be an ideal of R . Then $\sigma \in C_n(R, I)$ iff $\sigma_{ij}, \sigma_{ii} - \sigma_{jj} \in I$ for any $i \neq j$.

Definition

Let I be an ideal of R . Then the canonical map $R \rightarrow R/I$ induces a group homomorphism $\phi : GL_n(R) \rightarrow GL_n(R/I)$. The preimage of $Center(GL_n(R/I))$ under ϕ is called *full congruence subgroup of level I* and is denoted by $C_n(R, I)$.

Remark

Let $\sigma \in GL_n(R)$ and I be an ideal of R . Then $\sigma \in C_n(R, I)$ iff $\sigma_{ij}, \sigma_{ii} - \sigma_{jj} \in I$ for any $i \neq j$.

Theorem

Let I be an ideal of R . Then the Standard Commutator Formulas

$$[C_n(R, I), E_n(R)] = [E_n(R, I), E_n(R)] = E_n(R, I)$$

hold.

- 1 Introduction
- 2 The general linear group $GL_n(R)$
- 3 Sandwich classification for $GL_n(R)$**
- 4 The even-dimensional orthogonal group $O_{2n}(R)$
- 5 Sandwich classification for $O_{2n}(R)$
- 6 Sandwich classification for $U_{2n}(R, \Lambda)$
- 7 References

Definition

Let $\sigma \in GL_n(R)$. Then a matrix of the form $\epsilon \sigma^{\pm 1}$ where $\epsilon \in E_n(R)$ is called an *elementary σ -conjugate*.

Definition

Let $\sigma \in GL_n(R)$. Then a matrix of the form $\epsilon \sigma^{\pm 1}$ where $\epsilon \in E_n(R)$ is called an *elementary σ -conjugate*.

Theorem (RP, 2017)

Let $\sigma \in GL_n(R)$, $i \neq j$ and $k \neq l$. Then

- (i) $t_{kl}(\sigma_{ij})$ is a product of 8 elementary σ -conjugates and
- (ii) $t_{kl}(\sigma_{ii} - \sigma_{jj})$ is a product of 24 elementary σ -conjugates.

Definition

Let $\sigma \in GL_n(R)$. Then a matrix of the form ${}^\epsilon\sigma^{\pm 1}$ where $\epsilon \in E_n(R)$ is called an *elementary σ -conjugate*.

Theorem (RP, 2017)

Let $\sigma \in GL_n(R)$, $i \neq j$ and $k \neq l$. Then

- (i) $t_{kl}(\sigma_{ij})$ is a product of 8 elementary σ -conjugates and
- (ii) $t_{kl}(\sigma_{ii} - \sigma_{jj})$ is a product of 24 elementary σ -conjugates.

Proof.

(i) One checks easily that

$$[t_{12}(1), {}^{\tau^{-1}}[t_{32}(1), [\tau, \sigma]]] = t_{32}(\sigma_{23})$$

where $\tau = t_{21}(-\sigma_{23})t_{31}(\sigma_{22})$.

Definition

Let $\sigma \in GL_n(R)$. Then a matrix of the form ${}^\epsilon\sigma^{\pm 1}$ where $\epsilon \in E_n(R)$ is called an *elementary σ -conjugate*.

Theorem (RP, 2017)

Let $\sigma \in GL_n(R)$, $i \neq j$ and $k \neq l$. Then

- (i) $t_{kl}(\sigma_{ij})$ is a product of 8 elementary σ -conjugates and
- (ii) $t_{kl}(\sigma_{ii} - \sigma_{jj})$ is a product of 24 elementary σ -conjugates.

Proof.

(i) One checks easily that

$$[t_{12}(1), \tau^{-1}[t_{32}(1), [\tau, \sigma]]] = t_{32}(\sigma_{23})$$

where $\tau = t_{21}(-\sigma_{23})t_{31}(\sigma_{22})$. This implies that $t_{32}(\sigma_{23})$ is a product of 8 elementary σ -conjugates. Using the fact that $E_n(R)$ contains certain monomial matrices one can deduce (i).

(ii) Clearly the entry of $t_{ji}^{(1)}\sigma$ at position (j, i) equals $\sigma_{ii} - \sigma_{jj} + \sigma_{ji} - \sigma_{ij}$.

(ii) Clearly the entry of $t_{ji}^{(1)}\sigma$ at position (j, i) equals $\sigma_{ii} - \sigma_{jj} + \sigma_{ji} - \sigma_{ij}$. Applying (i) to $t_{ji}^{(1)}\sigma$ one gets that $t_{kl}(\sigma_{ii} - \sigma_{jj} + \sigma_{ji} - \sigma_{ij})$ is a product of 8 elementary σ -conjugates (clearly any elementary $t_{ji}^{(1)}\sigma$ -conjugate is also an elementary σ -conjugate).

(ii) Clearly the entry of $t_{ji}^{(1)}\sigma$ at position (j, i) equals $\sigma_{ii} - \sigma_{jj} + \sigma_{ji} - \sigma_{ij}$. Applying (i) to $t_{ji}^{(1)}\sigma$ one gets that $t_{kl}(\sigma_{ii} - \sigma_{jj} + \sigma_{ji} - \sigma_{ij})$ is a product of 8 elementary σ -conjugates (clearly any elementary $t_{ji}^{(1)}\sigma$ -conjugate is also an elementary σ -conjugate). Applying (i) to σ one gets that $t_{kl}(\sigma_{ij} - \sigma_{ji}) = t_{kl}(\sigma_{ij})t_{kl}(-\sigma_{ji})$ is a product of 16 elementary σ -conjugates.

(ii) Clearly the entry of $t_{ji}^{(1)}\sigma$ at position (j, i) equals $\sigma_{ii} - \sigma_{jj} + \sigma_{ji} - \sigma_{ij}$. Applying (i) to $t_{ji}^{(1)}\sigma$ one gets that $t_{kl}(\sigma_{ii} - \sigma_{jj} + \sigma_{ji} - \sigma_{ij})$ is a product of 8 elementary σ -conjugates (clearly any elementary $t_{ji}^{(1)}\sigma$ -conjugate is also an elementary σ -conjugate). Applying (i) to σ one gets that $t_{kl}(\sigma_{ij} - \sigma_{ji}) = t_{kl}(\sigma_{ij})t_{kl}(-\sigma_{ji})$ is a product of 16 elementary σ -conjugates. It follows that $t_{kl}(\sigma_{ii} - \sigma_{jj}) = t_{kl}(\sigma_{ii} - \sigma_{jj} + \sigma_{ji} - \sigma_{ij})t_{kl}(\sigma_{ij} - \sigma_{ji})$ is a product of 24 elementary σ -conjugates. □

Corollary

Let H be a subgroup of $GL_n(R)$. Then H is normalized by $E_n(R)$ iff

$$E_n(R, I) \subseteq H \subseteq C_n(R, I) \quad (1)$$

for some ideal I of R .

Corollary

Let H be a subgroup of $GL_n(R)$. Then H is normalized by $E_n(R)$ iff

$$E_n(R, I) \subseteq H \subseteq C_n(R, I) \quad (1)$$

for some ideal I of R .

Proof.

(\Rightarrow) Suppose that H is normalized by $E_n(R)$. Set $I = I(H)$. Then clearly $E_n(R, I) \subseteq H$. Let $\sigma \in H$, $i \neq j$ and $k \neq l$. By the previous theorem, $T_{kl}(\sigma_{ij})$ and $T_{kl}(\sigma_{ii} - \sigma_{jj})$ are products of matrices of the form ${}^\epsilon \sigma^{\pm 1}$ where $\epsilon \in E_n(R)$. Hence $T_{kl}(\sigma_{ij}), T_{kl}(\sigma_{ii} - \sigma_{jj}) \in H$ which implies $\sigma_{ij}, \sigma_{ii} - \sigma_{jj} \in I$. Thus $H \subseteq C_n(R, I)$.

Corollary

Let H be a subgroup of $GL_n(R)$. Then H is normalized by $E_n(R)$ iff

$$E_n(R, I) \subseteq H \subseteq C_n(R, I) \quad (1)$$

for some ideal I of R .

Proof.

(\Rightarrow) Suppose that H is normalized by $E_n(R)$. Set $I = I(H)$. Then clearly $E_n(R, I) \subseteq H$. Let $\sigma \in H$, $i \neq j$ and $k \neq l$. By the previous theorem, $T_{kl}(\sigma_{ij})$ and $T_{kl}(\sigma_{ii} - \sigma_{jj})$ are products of matrices of the form ${}^\epsilon \sigma^{\pm 1}$ where $\epsilon \in E_n(R)$. Hence $T_{kl}(\sigma_{ij}), T_{kl}(\sigma_{ii} - \sigma_{jj}) \in H$ which implies $\sigma_{ij}, \sigma_{ii} - \sigma_{jj} \in I$. Thus $H \subseteq C_n(R, I)$.

(\Leftarrow) Suppose that (1) holds for some ideal I . Then

$$[H, E_n(R)] \subseteq [C_n(R, I), E_n(R)] = E_n(R, I) \subseteq H.$$

Hence H is normalized by $E_n(R)$. □

- 1 Introduction
- 2 The general linear group $GL_n(R)$
- 3 Sandwich classification for $GL_n(R)$
- 4 The even-dimensional orthogonal group $O_{2n}(R)$**
- 5 Sandwich classification for $O_{2n}(R)$
- 6 Sandwich classification for $U_{2n}(R, \Lambda)$
- 7 References

Definition

Set $V := R^{2n}$. We consider V as right R -module and use the indexing $(e_1, \dots, e_n, e_{-n}, \dots, e_{-1})$ for the elements of the standard basis of V .

Definition

Set $V := R^{2n}$. We consider V as right R -module and use the indexing $(e_1, \dots, e_n, e_{-n}, \dots, e_{-1})$ for the elements of the standard basis of V . Define the quadratic form

$$q : V \rightarrow R$$
$$v \mapsto v^t \begin{pmatrix} 0 & p \\ 0 & 0 \end{pmatrix} v = \sum_{i=1}^n v_i v_{-i}$$

where $p \in M_n(R)$ is the matrix with ones on the skew diagonal and zeros elsewhere.

Definition

Set $V := R^{2n}$. We consider V as right R -module and use the indexing $(e_1, \dots, e_n, e_{-n}, \dots, e_{-1})$ for the elements of the standard basis of V . Define the quadratic form

$$q : V \rightarrow R$$
$$v \mapsto v^t \begin{pmatrix} 0 & p \\ 0 & 0 \end{pmatrix} v = \sum_{i=1}^n v_i v_{-i}$$

where $p \in M_n(R)$ is the matrix with ones on the skew diagonal and zeros elsewhere. The subgroup

$$O_{2n}(R) := \{ \sigma \in GL_{2n}(R) \mid q(\sigma v) = q(v) \ \forall v \in V \}$$

of $GL_{2n}(R)$ is called *(even-dimensional) orthogonal group*.

Lemma

Let $\sigma \in GL_{2n}(R)$. Then $\sigma \in O_{2n}(R)$ iff

- (i) $\sigma_{ij}^{-1} = \sigma_{-j,-i} \forall i, j \in \{1, \dots, n, -n, \dots, -1\}$ and
- (ii) $q(\sigma_{*j}) = 0 \forall j \in \{1, \dots, n, -n, \dots, -1\}$.

Lemma

Let $\sigma \in GL_{2n}(R)$. Then $\sigma \in O_{2n}(R)$ iff

- (i) $\sigma_{ij}^{-1} = \sigma_{-j,-i} \forall i, j \in \{1, \dots, n, -n, \dots, -1\}$ and
- (ii) $q(\sigma_{*j}) = 0 \forall j \in \{1, \dots, n, -n, \dots, -1\}$.

Lemma

Let $\sigma \in O_{2n}(R)$ and $k \in \{1, \dots, n, -n, \dots, -1\}$. Then the statements below are true.

- (i) If σ_{*k} is trivial then $\sigma_{-k,*}$ is trivial.
- (ii) If σ_{k*} is trivial then $\sigma_{*,-k}$ is trivial.

Definition

If $i, j \in \{1, \dots, n, -n, \dots, -1\}$ such that $i \neq \pm j$ and $x \in R$, then the matrix

$$T_{ij}(x) := e + xe^{ij} - xe^{-j,-i} \in O_{2n}(R)$$

is called an *elementary orthogonal transvection*. The subgroup of $O_{2n}(R)$ generated by all elementary orthogonal transvections is called *elementary orthogonal group* and is denoted by $EO_{2n}(R)$.

Definition

If $i, j \in \{1, \dots, n, -n, \dots, -1\}$ such that $i \neq \pm j$ and $x \in R$, then the matrix

$$T_{ij}(x) := e + xe^{ij} - xe^{-j,-i} \in O_{2n}(R)$$

is called an *elementary orthogonal transvection*. The subgroup of $O_{2n}(R)$ generated by all elementary orthogonal transvections is called *elementary orthogonal group* and is denoted by $EO_{2n}(R)$. Let I be an ideal of R . An elementary orthogonal transvection $T_{ij}(x)$ is called *I -elementary* iff $x \in I$. The subgroup of $O_{2n}(R)$ generated by all I -elementary orthogonal transvections is called *preelementary subgroup of level I* and is denoted by $EO_{2n}(I)$.

Definition

If $i, j \in \{1, \dots, n, -n, \dots, -1\}$ such that $i \neq \pm j$ and $x \in R$, then the matrix

$$T_{ij}(x) := e + xe^{ij} - xe^{-j,-i} \in O_{2n}(R)$$

is called an *elementary orthogonal transvection*. The subgroup of $O_{2n}(R)$ generated by all elementary orthogonal transvections is called *elementary orthogonal group* and is denoted by $EO_{2n}(R)$. Let I be an ideal of R . An elementary orthogonal transvection $T_{ij}(x)$ is called *I -elementary* iff $x \in I$. The subgroup of $O_{2n}(R)$ generated by all I -elementary orthogonal transvections is called *preelementary subgroup of level I* and is denoted by $EO_{2n}(I)$. The normal closure of $EO_{2n}(I)$ in $EO_{2n}(R)$ is called *elementary subgroup of level I* and is denoted by $EO_{2n}(R, I)$.

Lemma

The relations

$$T_{ij}(x) = T_{-j,-i}(-x), \quad (\text{R1})$$

$$T_{ij}(x)T_{ij}(y) = T_{ij}(x+y), \quad (\text{R2})$$

$$[T_{ij}(x), T_{hk}(y)] = e, \quad (\text{R3})$$

$$[T_{ij}(x), T_{jk}(y)] = T_{ik}(xy), \quad (\text{R4})$$

$$[T_{ij}(x), T_{j,-i}(y)] = e \quad (\text{R5})$$

hold where $h \neq j, -i$ and $k \neq i, -j$ in (R3) and $i \neq \pm k$ in (R4).

Definition

Let I be an ideal of R . Then the canonical map $R \rightarrow R/I$ induces a group homomorphism $\phi : O_{2n}(R) \rightarrow O_{2n}(R/I)$. The preimage of $\text{Center}(O_{2n}(R/I))$ under ϕ is called *full congruence subgroup of level I* and is denoted by $CO_{2n}(R, I)$.

Definition

Let I be an ideal of R . Then the canonical map $R \rightarrow R/I$ induces a group homomorphism $\phi : O_{2n}(R) \rightarrow O_{2n}(R/I)$. The preimage of $\text{Center}(O_{2n}(R/I))$ under ϕ is called *full congruence subgroup of level I* and is denoted by $CO_{2n}(R, I)$.

Remark

Let $\sigma \in O_{2n}(R)$ and I be an ideal of R . Then $\sigma \in CO_{2n}(R, I)$ iff $\sigma_{ij}, \sigma_{ii} - \sigma_{jj} \in I$ for any $i \neq j$.

Definition

Let I be an ideal of R . Then the canonical map $R \rightarrow R/I$ induces a group homomorphism $\phi : O_{2n}(R) \rightarrow O_{2n}(R/I)$. The preimage of $\text{Center}(O_{2n}(R/I))$ under ϕ is called *full congruence subgroup of level I* and is denoted by $CO_{2n}(R, I)$.

Remark

Let $\sigma \in O_{2n}(R)$ and I be an ideal of R . Then $\sigma \in CO_{2n}(R, I)$ iff $\sigma_{ij}, \sigma_{ii} - \sigma_{jj} \in I$ for any $i \neq j$.

Theorem

Let I be an ideal of R . Then the Standard Commutator Formulas

$$[CO_{2n}(R, I), EO_{2n}(R)] = [EO_{2n}(R, I), EO_{2n}(R)] = EO_{2n}(R, I)$$

hold.

- 1 Introduction
- 2 The general linear group $GL_n(R)$
- 3 Sandwich classification for $GL_n(R)$
- 4 The even-dimensional orthogonal group $O_{2n}(R)$
- 5 Sandwich classification for $O_{2n}(R)$**
- 6 Sandwich classification for $U_{2n}(R, \Lambda)$
- 7 References

Definition

Let $\sigma \in O_{2n}(R)$. Then a matrix of the form ${}^\epsilon\sigma^{\pm 1}$ where $\epsilon \in EO_{2n}(R)$ is called an *elementary orthogonal σ -conjugate*.

Definition

Let $\sigma \in O_{2n}(R)$. Then a matrix of the form ${}^\epsilon\sigma^{\pm 1}$ where $\epsilon \in EO_{2n}(R)$ is called an *elementary orthogonal σ -conjugate*.

Theorem (RP, 2017)

Let $\sigma \in O_{2n}(R)$, $i \neq \pm j$ and $k \neq \pm l$. Then

- (i) $T_{kl}(\sigma_{ij})$ is a product of 8 elementary orthogonal σ -conjugates,
- (ii) $T_{kl}(\sigma_{i,-i})$ is a product of 16 elementary orthogonal σ -conjugates,
- (iii) $T_{kl}(\sigma_{ii} - \sigma_{jj})$ is a product of 24 elementary orthogonal σ -conjugates and
- (iv) $T_{kl}(\sigma_{ii} - \sigma_{-i,-i})$ is a product of 48 elementary orthogonal σ -conjugates.

Definition

Let $\sigma \in O_{2n}(R)$. Then a matrix of the form ${}^\epsilon\sigma^{\pm 1}$ where $\epsilon \in EO_{2n}(R)$ is called an *elementary orthogonal σ -conjugate*.

Theorem (RP, 2017)

Let $\sigma \in O_{2n}(R)$, $i \neq \pm j$ and $k \neq \pm l$. Then

- (i) $T_{kl}(\sigma_{ij})$ is a product of 8 elementary orthogonal σ -conjugates,
- (ii) $T_{kl}(\sigma_{i,-i})$ is a product of 16 elementary orthogonal σ -conjugates,
- (iii) $T_{kl}(\sigma_{ii} - \sigma_{jj})$ is a product of 24 elementary orthogonal σ -conjugates and
- (iv) $T_{kl}(\sigma_{ii} - \sigma_{-i,-i})$ is a product of 48 elementary orthogonal σ -conjugates.

Proof.

$$[T_{12}(1), \tau^{-1} [T_{32}(1), [\tau, \sigma]]] = T_{32}(\sigma_{23})$$

where $\tau = T_{21}(-\sigma_{23})T_{31}(\sigma_{22})T_{2,-3}(\sigma_{2,-1})$. □

Corollary

Let H be a subgroup of $O_{2n}(R)$. Then H is normalized by $EO_{2n}(R)$ iff

$$EO_{2n}(R, I) \subseteq H \subseteq CO_{2n}(R, I)$$

for some ideal I of R .

- 1 Introduction
- 2 The general linear group $GL_n(R)$
- 3 Sandwich classification for $GL_n(R)$
- 4 The even-dimensional orthogonal group $O_{2n}(R)$
- 5 Sandwich classification for $O_{2n}(R)$
- 6 Sandwich classification for $U_{2n}(R, \Lambda)$**
- 7 References

Definition

Let $\sigma \in U_{2n}(R, \Lambda)$. Then a matrix of the form ${}^\epsilon \sigma^{\pm 1}$ where $\epsilon \in EU_{2n}(R, \Lambda)$ is called an *elementary unitary σ -conjugate*.

Definition

Let $\sigma \in U_{2n}(R, \Lambda)$. Then a matrix of the form ${}^\epsilon \sigma^{\pm 1}$ where $\epsilon \in EU_{2n}(R, \Lambda)$ is called an *elementary unitary σ -conjugate*.

Theorem (RP, 2017)

Let $\sigma \in U_{2n}(R, \Lambda)$, $k \neq \pm 1$ and $i \neq \pm j$. Then

- (i) $T_{kl}(\sigma_{ij})$ is a product of 160 elementary unitary σ -conjugates,
- (ii) $T_{kl}(\sigma_{i,-i})$ is a product of 320 elementary unitary σ -conjugates,
- (iii) $T_{kl}(\sigma_{ii} - \sigma_{jj})$ is a product of 480 elementary unitary σ -conjugates,
- (iv) $T_{kl}(\sigma_{ii} - \sigma_{-i,-i})$ is a product of 960 elementary unitary σ -conjugates and
- (v) $T_{k,-k}(\lambda^{-(\epsilon(k)+1)/2} |\sigma_{*j}|)$ is a product of $1600n + 4004$ elementary unitary σ -conjugates.

Definition

Let $\sigma \in U_{2n}(R, \Lambda)$. Then a matrix of the form ${}^\epsilon \sigma^{\pm 1}$ where $\epsilon \in EU_{2n}(R, \Lambda)$ is called an *elementary unitary σ -conjugate*.

Theorem (RP, 2017)

Let $\sigma \in U_{2n}(R, \Lambda)$, $k \neq \pm 1$ and $i \neq \pm j$. Then

- (i) $T_{kl}(\sigma_{ij})$ is a product of 160 elementary unitary σ -conjugates,
- (ii) $T_{kl}(\sigma_{i,-i})$ is a product of 320 elementary unitary σ -conjugates,
- (iii) $T_{kl}(\sigma_{ii} - \sigma_{jj})$ is a product of 480 elementary unitary σ -conjugates,
- (iv) $T_{kl}(\sigma_{ii} - \sigma_{-i,-i})$ is a product of 960 elementary unitary σ -conjugates and
- (v) $T_{k,-k}(\lambda^{-(\epsilon(k)+1)/2} |\sigma_{*j}|)$ is a product of $1600n + 4004$ elementary unitary σ -conjugates.

In the proof of the theorem above matrices like

$$\tau = T_{21}(\bar{\sigma}_{23}\sigma_{23})T_{31}(-\bar{\sigma}_{23}\sigma_{22})T_{3,-2}(\bar{\sigma}_{23}\sigma_{2,-1})T_{3,-3}(-\bar{\sigma}_{22}\sigma_{2,-1} + \bar{\lambda}\bar{\sigma}_{2,-1}\sigma_{22})$$

in the stabilizer of the second row of σ are used.

Corollary

Let H be a subgroup of $U_{2n}(R, \Lambda)$. Then H is normalized by $EU_{2n}(R, \Lambda)$ iff

$$EU_{2n}((R, \Lambda), (I, \Gamma)) \subseteq H \subseteq CU_{2n}((R, \Lambda), (I, \Gamma))$$

for some form ideal (I, Γ) of (R, Λ) .

- 1 Introduction
- 2 The general linear group $GL_n(R)$
- 3 Sandwich classification for $GL_n(R)$
- 4 The even-dimensional orthogonal group $O_{2n}(R)$
- 5 Sandwich classification for $O_{2n}(R)$
- 6 Sandwich classification for $U_{2n}(R, \Lambda)$
- 7 References

R. Preusser, *Sandwich classification for $GL_n(R)$, $O_{2n}(R)$ and $U_{2n}(R, \Lambda)$ revisited*,
accepted by J. Group Theory

R. Preusser, *Sandwich classification for $GL_n(R)$, $O_{2n}(R)$ and $U_{2n}(R, \Lambda)$ revisited*,
accepted by J. Group Theory

Thank you!